

ISSN 0104-6497

A stylized, high-contrast illustration of a palm tree. The trunk is a simple vertical line, and the fronds are depicted as a series of overlapping, curved shapes that create a sense of depth and texture. The illustration is rendered in a dark blue color against a lighter background.

NAUPLIUS

VOLUME 10

NUMBER 2

2002

NAUPLIUS

ISSN 0104-6497

THE JOURNAL OF THE BRAZILIAN CRUSTACEAN SOCIETY

Aims and Scope

The *Nauplius* publishes original articles based on research in any aspect of crustacean biology, including taxonomy, phylogeny, morphology, development, physiology, ecology, biogeography, bioenergetics, aquaculture and fisheries biology. Review articles with bold proposals showing original conclusions or approaches are encouraged.

The manuscripts submitted to this journal must have not been submitted to another periodical. Reproduction of articles published in the *Nauplius* is forbidden without previous written authorization by the Editorial Board of this periodical.

Instructions for the preparation of manuscripts are located on the inside back cover.

Manuscripts should be submitted to the Editor of *Nauplius*:

Maria Lucia Negreiros Fransozo
Departamento de Zoologia
Instituto de Biociências UNESP
18618-000, Botucatu, SP, Brasil

BRAZILIAN CRUSTACEAN SOCIETY

Officers:

Sérgio Luís de Siqueira Bueno (IB – USP, São Paulo, SP) President
Maria Lucia Negreiros Fransozo (IB – UNESP, Botucatu, SP) Secretary
Carlos Eduardo Falavigna da Rocha (IB – USP, São Paulo, SP) Treasurer
Adilson Fransozo (IB – UNESP, Botucatu, SP)
Fosca Pedini Pereira Leite (IB – UNICAMP, Campinas, SP)
Fernando Luís Medina Mantelatto (IB – USP, Ribeirão Preto, SP)

Brazilian Crustacean Society - SBC
Departamento de Zoologia - Instituto de Biociências - Universidade de São Paulo
Rua do Matão, Travessa 14 no.101
05508-900 São Paulo, SP, Brasil

Copyright by the Brazilian Crustacean Society

Summary

- **Population structure of *Temora stylifera* (Dana, 1849) (Copepoda) off Ipanema and in Guanabara Bay (Rio de Janeiro, Brazil).**
Schutze, M. L. M. and Björnberg, T. K. S. 0083-95
- **Spatial distribution of *Callicbirus major* (Say 1818) (Decapoda: Callianassidae) on a sandy beach, Piedade, Pernambuco, Brazil.**
Botter-Carvalho, M. L.; Santos, P. J. P. dos and Carvalho, P. V. V. da C. 097-109
- **Seasonal larval composition and abundance of shrimps in the surrounding area of the Patos Lagoon Mouth.**
Calazans, D. 111-120
- **Estuarine carcinofauna (Decapoda) of Rio Grande do Norte, Brazil.**
Ferreira, A. C. and Sankarankutty, C. 121-129
- **Growth patterns of *Portunus spinicarpus* (Stimpson, 1871) (Decapoda, Portunoidea) from Ubatuba region (SP), Brazil.**
Corbi-Corrêa, E. and Fransozo, A. 131-137
- **Cheliped morphometric differences in *Eurytium limosum* (Say, 1818) (Brachyura, Xanthidae).**
Guimarães, F. J. and Negreiros-Fransozo, M. L. 139-144
- **First record of *Pomatogebia operculata* (Schmitt, 1924) (Thalassinidea: Upogebiidae), in Venezuelan waters.**
Lira, C.; Blanco-Rambla, J. P.; Bolaños, J.; Hernández, G. and Hernández, J. 145-148
- **Description of reproductive performance and ovarian histology of wild *Farfantepenaeus paulensis* from shallow waters in southern Brazil.**
Peixoto, S.; Cavalli, R. O.; D'Incao, F.; Wasielesky, W. and Aguado, N. 149-153
- **Range extension of the geographic distribution of lobsters (Palinuroidea) and crabs (Xanthoidea) in the Brazilian coast.**
Cobo, V. J.; Pinheiro, A. P.; Freire, F. A. M. and Martins, I. A. 155-158